



# Targeting Phosphatidylserine with Calcium-Dependent Protein–Drug Conjugates for the Treatment of Cancer

Ran Li<sup>1</sup>, Srinivas Chiguru<sup>2</sup>, Li Li<sup>2</sup>, Dongyoung Kim<sup>3</sup>, Ramraj Velmurugan<sup>1,4</sup>, David Kim<sup>3</sup>, Siva Charan Devanaboyina<sup>1</sup>, Hong Tian<sup>5</sup>, Alan Schroit<sup>6</sup>, Ralph P. Mason<sup>2</sup>, Raimund J. Ober<sup>1,3</sup>, and E. Sally Ward<sup>1,7</sup>

## Abstract

In response to cellular stress, phosphatidylserine is exposed on the outer membrane leaflet of tumor blood vessels and cancer cells, motivating the development of phosphatidylserine-specific therapies. The generation of drug-conjugated phosphatidylserine-targeting agents represents an unexplored therapeutic approach, for which antitumor effects are critically dependent on efficient internalization and lysosomal delivery of the cytotoxic drug. In the current study, we have generated phosphatidylserine-targeting agents by fusing phosphatidylserine-binding domains to a human IgG1-derived Fc fragment. The tumor localization and pharmacokinetics of several phosphatidylserine-specific Fc fusions have been analyzed in mice and demonstrate that Fc-Syt1, a fusion containing the synaptotagmin 1 C2A domain,

effectively targets tumor tissue. Conjugation of Fc-Syt1 to the cytotoxic drug monomethyl auristatin E results in a protein–drug conjugate (PDC) that is internalized into target cells and, due to the  $\text{Ca}^{2+}$  dependence of phosphatidylserine binding, dissociates from phosphatidylserine in early endosomes. The released PDC is efficiently delivered to lysosomes and has potent antitumor effects in mouse xenograft tumor models. Interestingly, although an engineered, tetravalent Fc-Syt1 fusion shows increased binding to target cells, this higher avidity variant demonstrates reduced persistence and therapeutic effects compared with bivalent Fc-Syt1. Collectively, these studies show that finely tuned,  $\text{Ca}^{2+}$ -switched phosphatidylserine-targeting agents can be therapeutically efficacious. *Mol Cancer Ther*; 17(1): 169–82. ©2017 AACR.

## Introduction

The use of antibody-based targeted therapies for cancer has greatly expanded over the past two decades. However, almost all antibody-based agents target protein receptors that are present at higher levels on tumors compared with normal tissue. Although many tumor cells and the tumor vascular endothelium (1–3) specifically expose the negatively charged phospholipid, phosphatidylserine, in their outer membrane leaflet, few studies have taken advantage of phosphatidylserine as a therapeutic target (4).

The identification of phosphatidylserine as a tumor marker has prompted the development of molecular targeted therapies specific for this phospholipid. Among these therapeutics is bavitux-

imab, a chimeric mAb that targets the phosphatidylserine-binding serum protein  $\beta 2$  glycoprotein-1 ( $\beta 2\text{GP1}$ ; ref. 5). By cross-linking  $\beta 2\text{GP1}$ , bavituximab binds phosphatidylserine indirectly with high affinity and elicits antiangiogenic effects via antibody-dependent cell-mediated cytotoxicity (6). In addition, by masking phosphatidylserine in the tumor microenvironment, bavituximab might also play an immunomodulatory role by inducing polarization of macrophages to the inflammatory M1 phenotype and reducing the number of myeloid-derived suppressor cells (7). In combination with radiation or chemotherapy, bavituximab has been demonstrated to be effective in multiple preclinical models for glioblastoma, pancreatic cancer, prostate, breast, and hepatocellular carcinomas (6–10). However, a recent phase III clinical trial (NCT01999673; <http://www.peregrineinc.com/pipeline/bavituximab-oncology.html>) in non-small cell lung cancer patients was discontinued as bavituximab in combination with docetaxel did not show sufficient improvement in overall survival over treatment with docetaxel alone. This motivates the development of strategies to improve the *in vivo* antitumor activity of phosphatidylserine-targeting therapies.

Antibody–drug conjugates (ADCs) combine the high specificity of antibody targeting with potent cytotoxic drugs and promise to be more effective in killing tumor cells than their corresponding "naked" antibodies (11, 12). To date, three ADCs (Mylotarg, Adcetris, and Kadcyla) have received regulatory approval, although Mylotarg has been withdrawn (11). More than 50 other ADCs are in clinical development (13, 14). This suggests that a phosphatidylserine-specific ADC may have improved efficacy over naked phosphatidylserine-targeting antibodies, such as bavituximab. However, the generation of a phosphatidylserine-

<sup>1</sup>Department of Molecular and Cellular Medicine, Texas A&M University Health Science Center, College Station, Texas. <sup>2</sup>Department of Radiology, University of Texas Southwestern Medical Center, Dallas, Texas. <sup>3</sup>Department of Biomedical Engineering, Texas A&M University, College Station, Texas. <sup>4</sup>Biomedical Engineering Graduate Program, University of Texas Southwestern Medical Center, Dallas, Texas. <sup>5</sup>China Pharmaceutical University, Nanjing, Jiangsu, China. <sup>6</sup>Department of Pharmacology, University of Texas Southwestern Medical Center, Dallas, Texas. <sup>7</sup>Department of Microbial Pathogenesis and Immunology, Texas A&M University Health Science Center, College Station, Texas.

**Note:** Supplementary data for this article are available at Molecular Cancer Therapeutics Online (<http://mct.aacrjournals.org/>).

**Corresponding Author:** E. Sally Ward, Department of Molecular and Cellular Medicine, Texas A&M University Health Science Center, College Station, TX 77843. Phone: 979-436-0742; Fax: 979-847-9481; E-mail: [sally.ward@tamu.edu](mailto:sally.ward@tamu.edu)

doi: 10.1158/1535-7163.MCT-17-0092

©2017 American Association for Cancer Research.

specific ADC faces several challenges. First, following binding to the cell surface, ADCs require effective internalization into the lysosomal pathway. The internalization behavior of phosphatidylserine and its subsequent trafficking have not been explored. Second, in contrast with protein antigens, the development of high-affinity, specific antibodies to small-molecule "haptens" such as phosphatidylserine is challenging (bavituximab targets phosphatidylserine indirectly through binding to  $\beta$ 2GP1). Third, under certain circumstances phosphatidylserine can also be exposed on the surface of nonapoptotic cells, such as differentiating monocytes and a subpopulation of T cells (15–17).

To generate an effective phosphatidylserine-directed drug conjugate, we produced a panel of phosphatidylserine-targeting agents by fusing naturally occurring phosphatidylserine-binding domains to the Fc portion of human IgG1. To achieve efficient dissociation of the targeting agents from phosphatidylserine in sorting (early) endosomes, followed by their lysosomal delivery, we have exploited the significant decrease in  $\text{Ca}^{2+}$  levels in endosomes (18, 19) by using phosphatidylserine-binding domains that interact with phosphatidylserine in a  $\text{Ca}^{2+}$ -dependent manner. We initially analyzed the pharmacokinetics and tumor targeting of the phosphatidylserine-targeting agents in mice, and observed that despite its relatively low affinity for phosphatidylserine binding, a fusion comprising the C2A domain of synaptotagmin 1 (Fc-Syt1) had superior properties compared with higher affinity protein kinase C- $\alpha$  (PKC $\alpha$ )- and Annexin A1 (AnxA1)-based fusions. Fc-Syt1 was therefore used as a platform to generate a protein–drug conjugate (PDC) with monomethyl auristatin E (MMAE). Modulation of the avidity of Fc-Syt1 for phosphatidylserine binding demonstrated that a bivalent Fc-Syt1-based PDC had improved therapeutic efficacy over a tetravalent variant due to more favorable pharmacokinetic properties. Our observations indicate the need for affinity/avidity tuning of phosphatidylserine-targeting PDCs and provide novel insight into the design of therapeutics to target this phospholipid.

## Materials and Methods

### Cell lines and culture conditions

2H11 mouse tumor endothelial cells (ATCC, CRL-2163) and MDA-MB-231 human breast cancer cells (ATCC, HTB-26) were cultured in DMEM supplemented with 5% and 10% FBS, respectively. T-47D breast cancer cells (ATCC, HTB-133), LNCaP and 22Rv1 human prostate cancer cells (ATCC, CRL1740 and CRL-2505) were cultured in RPMI1640 medium supplemented with 10% FBS. SK-BR-3 human breast cancer cells (ATCC, HTB-30) were cultured in McCoy's 5A medium supplemented with 10% FBS. All cell lines were authenticated with DNA fingerprinting by the University of Arizona Genetics Core on November 16, 2016, and tested for mycoplasma at monthly intervals. All cell lines were used within 15 passages following thawing of master stocks. Cells were cultured at 37°C with 5%  $\text{CO}_2$ . Expi293F cells were used for protein expression and were cultured in Expi293 expression medium (Life Technologies, catalog number A14635) at 37°C with 8%  $\text{CO}_2$  and 80% humidity.

### Antibodies

The following antibodies were used in this study: rat anti-mouse LAMP1, mouse anti-human LAMP1, and mouse anti- $\beta$ -tubulin antibodies (Developmental Studies Hybridoma Bank, clone # 1D4B, H4A3 and E7); mouse anti-human EEA1 and rat

anti-mouse CD31 antibodies (BD Biosciences, catalog # 610456 and 557355); goat anti-human IgG (H + L) antibody conjugated with HRP, donkey anti-rat (H + L) antibody conjugated with Alexa Fluor 488 and donkey anti-human IgG (H + L) antibody conjugated with Cy3 (Jackson ImmunoResearch, catalog # 109-035-003, 712-545-153 and 709-165-149); goat anti-human IgG (H + L) antibody conjugated with Alexa Fluor 555, goat anti-mouse IgG (H + L) antibody conjugated with Alexa Fluor 488 and goat anti-human IgG (H + L) antibody conjugated with Alexa Fluor 647 (Life Technologies, catalog # A21433, A11029, and A21445); rabbit anti-human Ki-67 antibody (Abcam, catalog # 92742).

### Protein labeling

Proteins were labeled with Alexa 647 or IRDye800CW with the Alexa Fluor 647 NHS Ester (Life Technologies, catalog # A37573) or IRDye800CW Protein Labeling Kit (LI-COR, catalog # 928-38040), respectively, following the manufacturer's instructions. Proteins were labeled with I-125 as described previously (20). Radio-TLC analysis was used to determine radioiodination efficiency and radiochemical purity using a Rita Star Radioisotope TLC Analyzer with ITLC-SG plates and PBS as the mobile phase. The typical radiochemical yield was 68% to 76%, with radiochemical purity of >99.5% and specific activity of 10 to 12  $\mu\text{Ci}/\mu\text{g}$ . The absence of aggregation for fluorophore-labeled proteins was verified by gel filtration analyses.

### Generation of expression constructs

For use as controls, the Fc region including the hinge region (residues 214–446 of the heavy chain) of the hen egg lysozyme-specific human IgG1, HuLys10 (21), was cloned into the pcDNA3.4 vector (Invitrogen, catalog # 14308) with the N-terminal leader peptide derived from a mouse IgG heavy chain (21, 22). Similarly, the genes encoding the heavy and light chain genes (cDNA) of the HuLys10 antibody were cloned into pcDNA3.4. For the control IgG heavy and light chain constructs, Cys214 in the light chain, which forms a sulfhydryl bridge with Cys219 in the heavy chain of HuLys10 were both mutated to serine residues using the QuikChange II Site-Directed Mutagenesis Kit (Agilent Technologies, catalog # 200523).

cDNA clones for human Annexin A1 (AnxA1), human synaptotagmin 1 (Syt1), and human PKC $\alpha$  were purchased from Open Biosystems (clone ID: 3459615, clone ID: 6187902 and clone ID: 40028305, respectively). Genes encoding the AnxA1 phosphatidylserine-binding core domain (amino acids 41–346), the Syt1 phosphatidylserine-binding C2A domain (amino acids 141–266) and PKC $\alpha$  phosphatidylserine-binding C2 domain (amino acids 157–288) were fused via a Gly<sub>4</sub>Ser linker sequence to the CH<sub>3</sub> domain of the human IgG1 Fc region (residues 214–446) with a leader peptide derived from the mouse IgG heavy chain (21, 22). Cys219 in the hinge region was mutated in all Fc fusion constructs so that there are two cysteine residues per hinge. The genes encoding the Fc fusions were cloned into the pcDNA3.4 vector.

To generate Syt1-Fc-Syt1, the Syt1 phosphatidylserine-binding C2A domain (amino acids 141–266) was linked to the N-terminus of the hinge region of the Fc-Syt1 construct via a Gly<sub>4</sub>Ser linker sequence. The leader peptide derived from a mouse IgG heavy chain (21, 22) was appended to the N-terminus of the hinge-linked Syt1 phosphatidylserine-binding C2A domain, and the resulting Fc fusion was cloned into the pcDNA3.4 vector.

Mutations to reduce phosphatidylserine binding of the Syt1 C2A domain (D173N, D179N, D231N, D233N, and D239N; ref. 23) were inserted in the Fc-Syt1 construct to generate Fc-Syt1 (DN) and cloned into the pcDNA3.4 vector. All constructs were generated using standard methods of molecular biology and designed oligonucleotides. Following construction of expression plasmids, sequences were verified. Oligonucleotide and construct sequences are available upon request.

#### Protein expression and purification

All phosphatidylserine-targeting proteins used in this study were produced using the Expi293 expression system from Life Technologies following the manufacturer's instructions. Briefly, cells were transfected with expression constructs for 6 to 7 days and Fc fusion proteins were purified from culture supernatants using protein G-Sepharose. Bound proteins were eluted using 50 mmol/L diethylamine with 150 mmol/L NaCl. The eluted protein was neutralized using 2 mol/L Tris pH 7.0 followed by dialysis against PBS. All proteins were concentrated and loaded onto a Hiload 16/60 Superdex 200 gel filtration column (GE Healthcare). The homodimeric, nonaggregated form of the protein was separated, concentrated, and analyzed using a Superdex 200 15/30 gel filtration column (GE Healthcare).

Recombinant mouse FcRn was produced as described previously (24). Briefly, High Five cells grown at 27°C in EX-CELL 405 medium (Sigma, catalog # 14405) were infected at a density of  $1 \times 10^6$  cells/mL with recombinant baculovirus (mouse FcRn  $\alpha$ -chain/mouse  $\beta$ 2-microglobulin). Cells were cultured at 23°C to 24°C for 72 hours. Mouse FcRn was purified from the supernatant using  $\text{Ni}^{2+}$ -NTA agarose (Qiagen) followed by the use of a Hiload 16/60 Superdex 200 gel filtration column.

#### MALDI-TOF mass spectrometry

For MALDI-TOF analyses, all samples (20–40  $\mu\text{g}$  diluted in 20  $\mu\text{L}$  PBS) were treated with 1  $\mu\text{L}$  (10 U) PNGase F (Promega, catalog # V4831) at 37°C for 1 hour. Samples were extracted using C<sub>4</sub> LithTips (Protea, catalog # SP-410) and analyzed with sinapic acid under linear conditions with a Kratos/Shimadzu Axima CFR MALDI-TOF mass spectrometer.

#### Surface plasmon resonance (BIAcore) analyses

Surface plasmon resonance experiments were carried out using a BIAcore T200 (GE Healthcare). To determine the equilibrium binding affinities of mouse FcRn for the phosphatidylserine-targeting agents (Fc-AnxA1, Fc-Syt1, and Fc-PKC $\alpha$ ), control human IgG1 and Fc, mouse FcRn was injected over immobilized proteins (coupled at ~750–1,500 RU on flow cells of CM5 sensor chips) at a flow rate of 10  $\mu\text{L}/\text{minute}$  in PBS (pH 6.0) with 0.01% v/v Tween 20. Flow cells were regenerated at the end of each run using 0.15 mol/L NaCl, 0.1 mol/L  $\text{NaHCO}_3$ , pH 8.5. The dissociation constants ( $K_D$ s) for the interactions of mouse FcRn with different proteins were determined using custom written software (25, 26). IgG or Fc has two possible interaction sites on mouse FcRn, and equilibrium binding data were fitted as described previously (26). The equilibrium dissociation constants for the higher affinity interaction site are reported.

#### Membrane lipid strip binding assay

Lipid-coated membrane strips (Echelon, catalog # P-6002) were first hydrated with TBST (20 mmol/L Tris, 150 mmol/L NaCl, 0.1% Tween 20, pH 7.5) and then incubated with blocking solution (4% fatty acid free BSA dissolved in TBST) at room

temperature for 1 hour. Proteins were diluted to a concentration of 2  $\mu\text{g}/\text{mL}$  in blocking buffer and incubated with membranes at room temperature for 2 hours. The lipid strip was then washed with TBST and bound proteins detected using HRP-conjugated goat anti-human IgG (H + L) antibody.

#### Fluorescence microscopy analyses

2H11 or MDA-MB-231 cells were grown on coverslips (Zeiss, ref # 0109030091) and incubated with either control human IgG1 (HuLys10) or phosphatidylserine-specific agents diluted in growth medium for the indicated time periods in figure legends. Cells were then washed with PBS and fixed with ice-cold 4% PFA at room temperature for 20 minutes. Following fixation, cells were permeabilized with 0.1% Triton X-100 and incubated with blocking buffer (PBS, 0.1% Tween 20, 5% serum) at room temperature for 30 minutes. Primary antibodies specific for mouse LAMP-1 (clone 1D4B), human LAMP-1 (clone H4A3), or mouse early endosomal antigen-1 (EEA1) were diluted in blocking buffer and incubated with cells at room temperature for 2 hours. For staining of microtubule networks, cells were fixed with methanol at  $-20^\circ\text{C}$  for 3 minutes. Following fixation, cells were incubated with blocking buffer (PBS, 0.1% Tween 20, 5% serum) at room temperature for 30 minutes. The primary antibody specific for  $\beta$ -tubulin was diluted in blocking buffer and incubated with cells at room temperature for 2 hours. After incubation with the primary antibodies, cells were washed with PBST (PBS, 0.1% Tween 20) and incubated with fluorescently labeled secondary antibodies diluted in blocking buffer for 1 hour at room temperature. To detect internalized phosphatidylserine-specific agents, fluorescently labeled goat or donkey anti-human IgG (H + L) antibody was used. Following incubation, cells were washed with PBST and mounted with ProLong Gold Antifade Mountant (Life Technologies, catalog # P36930).

Fluorescent images were acquired using a Zeiss Axiovert 200M inverted fluorescence microscope with a 63 $\times$ , 1.4 NA plan apochromat objective (Carl Zeiss) and a 1.6 $\times$  internal optovar. Fluorescent filter sets for GFP and Alexa 555 (Chroma Technology, catalog # 41017 and 41002b) were used for image acquisition. The acquired data were processed using the microscopy image analysis tool (MIATool) software ([www4.utsouthwestern.edu/wardlab/miatool.asp](http://www4.utsouthwestern.edu/wardlab/miatool.asp)).

#### Phosphatidylserine pull-down assay

To study the  $\text{Ca}^{2+}$ -dependence of binding of the phosphatidylserine-specific Fc fusions to phosphatidylserine, proteins (100 nmol/L) were diluted in binding buffer (10 mmol/L HEPES pH 7.4, 150 mmol/L NaCl with 2 mmol/L or 2  $\mu\text{mol}/\text{L}$   $\text{Ca}^{2+}$ ). Phosphatidylserine-coated beads (50  $\mu\text{L}$ ; Echelon, catalog # P-BOPS) were added and incubated at room temperature for 2 hours. The beads were then washed with the binding buffer and bound proteins detected by immunoblotting with HRP-conjugated goat anti-human IgG (H + L) antibody. To examine the pH-dependent binding to phosphatidylserine, proteins were diluted at 100 nmol/L in PBS pH 7.4 or 6.0. Fifty microliters (bed volume) phosphatidylserine-coated beads were added, incubated at room temperature for 2 hours, and samples processed as above.

#### Annexin V-binding assay

Cells ( $1 \times 10^6$ ) were suspended in Annexin V-binding solution (10 mmol/L HEPES pH 7.4, 150 mmol/L NaCl, 2.5 mmol/L  $\text{CaCl}_2$ ). Annexin V conjugated with Alexa 488 (Life Technologies,

catalog # A13201) was added to the cell suspension at a 1:100 dilution and incubated with the cells for 10 minutes at room temperature. Cells were then washed once with Annexin V-binding solution and analyzed by flow cytometry (BD FACS-Calibur). Flow cytometry data were processed using FlowJo (FlowJo, LLC).

#### Flow cytometry analyses of phosphatidylserine-targeting agents

Cells were trypsinized and resuspended in flow cytometry buffer (PBS with  $\text{Ca}^{2+}$ /  $\text{Mg}^{2+}$ , 1% BSA). Phosphatidylserine-specific Fc fusions (50 nmol/L) were incubated with the cells for 30 minutes at either room temperature or on ice depending on the assay. Cells were washed with flow cytometry buffer and incubated with fluorophore-conjugated secondary antibodies on ice for 30 minutes. Cells were then washed and analyzed by flow cytometry (BD FACS-Calibur). Flow cytometry data were processed using FlowJo (FlowJo LLC).

#### Protein conjugation with maleimidocaproyl-val-cit-PAB-MMAE

Fc fusion or control antibody (20  $\mu\text{mol/L}$ ) in PBS was incubated with 320  $\mu\text{mol/L}$  TCEP at room temperature for 3 hours to reduce the hinge disulfide bonds. Maleimidocaproyl-val-cit-PAB-MMAE (MC-VC-PAB-MMAE; Levena Biopharma, catalog # SET0201) was then added to the reduced Fc fusion or control antibody at a concentration of 160  $\mu\text{mol/L}$  and incubated at room temperature for 3 hours. Following the conjugation reaction, free MMAE was removed by dialysis of the protein against PBS. The conjugated Fc fusion or control antibody was stored at 4°C.

#### Cell growth and survival assays

Cells were plated into 96-well plates as follows: 2H11, 10,000 cells/well; MCF-7, SK-BR-3 and LNCaP, 5,000 cells/well; MDA-MB-231, 3,000 cells/well; 22Rv1, 15,000 cells/well. Cells were grown overnight followed by the addition of phosphatidylserine-specific PDCs. Cell growth and survival were measured after a 3- to 5-day incubation with the Cell Proliferation AQ One Solution Cell Proliferation Assay Kit (Promega, catalog # G3581). Dose-response curves were plotted and  $\text{IC}_{50}$  values determined using GraphPad Prism software.

#### Pharmacokinetic and whole-body imaging studies

Animal procedures used in all mouse studies were approved by the Institutional Animal Care and Use Committee of the University of Texas Southwestern Medical Center (Dallas, TX) and Texas A&M University (College Station, TX). All BALB/c SCID mice used in the study were purchased from The Jackson Laboratory (stock # 001803) and bred in-house. Pharmacokinetic studies were performed as described previously (20). Briefly, Lugol solution was added to drinking water 96 hours before the experiments. SCID BALB/c female mice (8 weeks old; 18–22 g weight) were anesthetized using 2% isoflurane in oxygen and injected (intravenously) with  $^{125}\text{I}$ -labeled proteins (100–120  $\mu\text{Ci}$ , 10–12  $\mu\text{g}/\text{mouse}$ ) in 200  $\mu\text{L}$  PBS with 0.1% BSA. Whole-body radioactive counts were measured using a Biodex Atomlab 100 dose calibrator. Blood radioactive counts were measured by retro-orbitally bleeding mice with 10  $\mu\text{L}$  capillary tubes (Drummond) and radioactive counts (cpm) determined by gamma counting (PerkinElmer).

For whole-body, near-infrared imaging (NIR), female nude mice (6–7 weeks old; purchased from Envigo, catalog # 6903F)

or BALB/c SCID mice (6–8 weeks old) were used. For implantation of MDA-MB-231 tumors, mice were anesthetized with 2% isoflurane in oxygen, and a small surgical incision was made to expose the mammary fat pad. MDA-MB-231 cells were trypsinized and dispersed into single-cell suspensions in PBS. A total of  $5 \times 10^6$  cells per mouse were injected in 100  $\mu\text{L}$  into the mammary fat pad, using a 25 G needle, and the incision was then sealed with a wound clip. Buprenorphine was administered (subcutaneously) at 50  $\mu\text{g}/\text{kg}$  immediately following the surgery and 12 hours later. Mice were monitored daily, and the wound clip was removed one week postsurgery. For imaging of nude mice, mice were divided into 3 groups ( $n = 3$  mice/group) when tumors reached sizes of approximately 150  $\text{mm}^3$  and administered (intraperitoneally) with 5 mg/kg docetaxel 72 and 48 hours before being injected (intravenously) with 1 nmol IRDye800CW-labeled phosphatidylserine-specific agents in PBS. Fluorescence imaging (FLI) was performed using a Caliper Xenogen IVIS Spectrum (PerkinElmer) *in vivo* imaging system at 0 (before injection) and 3, 24, and 48 hours postinjection. FLI was performed using 745 nm excitation, 800 nm emission, binning 8, FOV 12.9 cm, f-stop 2, and auto-exposure. Data were quantitated with the Living Imaging software using absolute radiant efficiency (photons/second) in an ROI, manually drawn to outline the FLI signal of the tumor and normalized to the tumor volumes.


For analyses of tumor localization in BALB/c SCID mice, mice were divided into three groups ( $n = 3$  mice/group) when tumors reached sizes of approximately 300  $\text{mm}^3$  and administered (intraperitoneally) with 5 mg/kg docetaxel 72 and 48 hours before being injected (intravenously) with 1 nmol of IRDye800CW-labeled phosphatidylserine-specific agents in PBS. Forty-eight hours following injection, tumors were dissected out and imaged as above. Fluorescence in an ROI, manually drawn to outline the FLI signal of the tumor, was quantitated and normalized to the tumor weight.

#### Mouse xenograft studies

Implantation of MDA-MBA-231 tumor xenografts in BALB/c SCID mice was carried out as described for the whole-body imaging experiments. For implantation of LNCaP tumors, 7- to 8-week-old male BALB/c SCID mice (18–22 g weight) were anesthetized with 2% isoflurane in oxygen and  $5 \times 10^6$  LNCaP cells suspended in 50% RPMI and 50% Matrigel (BD Biosciences) were injected (subcutaneously). When MDA-MBA-231 or LNCaP tumors reached a size of approximately 100  $\text{mm}^3$ , mice were injected (intraperitoneally) with 5 mg/kg docetaxel 72 and 48 hours before the treatment. Mice were then injected (intravenously) with 1 nmol unconjugated proteins, PDCs, or PBS vehicle as indicated in the figure legends twice per week. Tumors and body weights were measured twice a week. For the treatment experiment using Fc-Syt1(DN)\_MMAE, mice were treated for 4 weeks and monitored for another 2.5 weeks. Experiments were terminated when the tumor size reached 2 cm in any dimension.

#### IHC analyses

Female BALB/c SCID mice bearing MDA-MB-231 tumors were treated (intraperitoneally) with 5 mg/kg docetaxel 72 and 48 hours before intravenous delivery of either PBS vehicle or 1 nmol Fc-Syt1 conjugated with MMAE. At different time points, mice were perfused (intracardiac) with PBS followed by 4% PFA. Tumors were then dissected out, embedded in OCT (Thermo Fisher Scientific, catalog # 23-730-571) and stored at  $-80^\circ\text{C}$ . Ten-

**Figure 1.**

Generation and characterization of phosphatidylserine-targeting agents. **A**, Schematic representation of phosphatidylserine agents (left). Filled circles and rectangles represent the phosphatidylserine-binding domains and IgG1 hinge region, respectively. Right, reducing SDS-PAGE analyses of the phosphatidylserine-specific Fc fusions, with molecular weights (MW) shown in kDa on the left. **B**, Lipid-binding profiles of phosphatidylserine-specific Fc fusions using lipid-coated nitrocellulose membranes. Bound proteins were detected with goat anti-human IgG (H + L) antibody conjugated with HRP. **C**, Binding of phosphatidylserine-specific Fc fusions to phosphatidylserine-positive 2H11 and MDA-MB-231 cells using flow cytometry analysis, using Alexa 647-labeled anti-human IgG (H + L) for detection. 2<sup>nd</sup> only and Fc represent negative controls using secondary conjugate or recombinant Fc fragment, respectively. **D**, Pharmacokinetic analyses of phosphatidylserine-specific Fc fusions in BALB/c SCID mice ( $n = 5$  mice/group). Whole body and blood levels of radioactivity were measured at the indicated time points. **E**, AUCs in **D** for whole-body (top) and blood (bottom) counts were quantitated. **F**, Nude mice bearing orthotopic human MDA-MB-231 tumors ( $n = 3$  mice/group) were injected with IRDye800CW-labeled phosphatidylserine-specific Fc fusions, and NIR fluorescence images acquired at the indicated time points. **G**, Tumor-associated fluorescence intensities at 48 hours in **F** normalized to the corresponding tumor volumes were quantitated. **H**, Female BALB/c SCID mice bearing MDA-MB-231 tumors ( $n = 3$  mice/group) were injected with IRDye800CW-labeled phosphatidylserine-specific agents. Forty-eight hours postinjection, tumors were dissected out and NIR images were acquired. **I**, Tumor-associated fluorescence intensities in **H** normalized to the corresponding tumor weights. Statistically significant differences in **E**, **G**, and **I** were analyzed using one-way ANOVA followed by Tukey *post hoc* test (\*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ ; \*\*\*\*,  $P < 0.0001$ ). Error bars in **D**, **E**, **G**, and **I** represent SEM.

micron tissue sections were cut and hydrated with PBS at room temperature before fixation with 4% PFA for 30 minutes. Tumor sections were then washed with PBS and incubated with the permeabilization/blocking solution (PBS, 0.5% Triton X-100, 3% BSA) at room temperature for 1 hour. Primary antibodies for human Ki-67, mouse CD31, and F4/80 were diluted in the blocking buffer (PBS, 0.1% Tween 20, 5% serum) and incubated with the tissue sections at 4°C overnight. The next day, tissue sections were washed with PBST (PBS, 0.1% Tween 20) and incubated with the fluorophore-conjugated secondary antibodies diluted in the blocking buffer at room temperature for 2 hours. After washing with PBST, tissue sections were mounted with ProLong Gold antifade mounting medium. Confocal images were acquired using a Nikon A1R confocal microscope equipped with a 40×, 1.3 NA plan fluor objective and processed with the NIS-Elements software (Nikon).

## Results

### Generation and characterization of phosphatidylserine-targeting agents

A panel of phosphatidylserine-targeting agents was generated by fusing the Fc region of human IgG1 to the following phosphatidylserine-binding domains: core domain of Annexin A1 (AnxA1), C2A domain of Synaptotagmin 1 (Syt1), and C2 domain of PKCα (Fig. 1A). The three domains bind to phosphatidylserine in a Ca<sup>2+</sup>-dependent manner (4, 27, 28), and the resulting fusion proteins were designated Fc-AnxA1, Fc-Syt1, and Fc-PKCα, respectively. The phosphatidylserine-targeting agents were expressed and purified as homodimers (Fig. 1A; Supplementary Fig. S1) and bound to phosphatidylserine in a lipid-binding assay (Fig. 1B). They also bound to cardiolipin, which is located on the inner mitochondrial membrane of eukaryotic cells and therefore not relevant to targeting. In contrast with Fc-Syt1 and Fc-PKCα, Fc-AnxA1 exhibited a broader lipid-binding profile and bound both neutral and negatively charged lipids (Fig. 1B). Importantly, none of the phosphatidylserine-binding agents bound phosphatidylcholine and sphingomyelin, lipids that are present in the plasma membrane outer leaflet.

The tumor endothelial cell line 2H11 (29) and triple-negative breast tumor cell line MDA-MB-231 were used to investigate the ability of the phosphatidylserine-binding agents to interact with lipids on the cell surface. Binding of fluorescent Annexin V showed that both of these cell lines expose phosphatidylserine and that phosphatidylserine exposure increases following docetaxel treatment (Supplementary Fig. S2). All phosphatidylserine-binding agents interacted with phosphatidylserine-positive cells, with Fc-Syt1 showing lower levels of binding than Fc-AnxA1 and Fc-PKCα (Fig. 1C).

**Table 1.** Equilibrium dissociation constants for the interactions of recombinant proteins with mouse FcRn at pH 6.0 and 7.4 determined using surface plasmon resonance

Proteins	K <sub>D</sub> (nmol/L), pH 6.0	K <sub>D</sub> (nmol/L), pH 7.4
Human IgG1 <sup>a</sup>	57.1	n.b. <sup>b</sup>
Fc	66.8	n.b. <sup>b</sup>
Fc-AnxA1	74.1	n.b. <sup>b</sup>
Fc-Syt1	69.4	n.b. <sup>b</sup>
Fc-PKCα	69.1	n.b. <sup>b</sup>

<sup>a</sup>Human IgG1 is the hen egg lysozyme-specific antibody, HuLys10 (21).

<sup>b</sup>n.b., no detectable binding, or affinity too low to accurately determine.

To determine which recombinant proteins were suitable for further development as PDCs, we evaluated the pharmacokinetic behavior and tumor localization of the three phosphatidylserine-binding agents in mice. Pharmacokinetic studies of the phosphatidylserine-binding agents revealed that Fc-Syt1 persisted at both the whole-body and blood levels for significantly longer than the other two proteins (Fig. 1D and E). Binding analyses demonstrated that these three fusion proteins had similar affinities for binding to recombinant mouse FcRn at pH 6.0, with no detectable binding or affinity too low to accurately determine at pH 7.4 (Table 1), indicating that affinity differences do not contribute to the longer persistence of Fc-Syt1. The proteins were labeled with the residualizing dye, IRDye800CW, and injected (intravenously) into female nude mice bearing MDA-MB-231 xenografts and imaged at different time points (Fig. 1F) with tumor fluorescence quantitated 48 hours postinjection (Fig. 1G). Similar experiments were carried out in BALB/c SCID mice, and 48 hours later, tumors were excised and dye levels determined (Fig. 1H and I). Among the three phosphatidylserine-specific agents, Fc-Syt1 exhibited the highest level of tumor localization. Collectively, these results indicated that of the three phosphatidylserine-targeting agents, Fc-Syt1 is the most promising for conversion to a PDC for use in therapy.

### Cell binding and internalization of bivalent and tetravalent phosphatidylserine-specific Fc fusions

For therapeutic efficacy of an antibody–drug conjugate (ADC), target cell internalization of the ADC and release of the toxic payload in late endosomes or lysosomes is essential (11). In several receptor systems, it has been shown that multivalent ligands, or mixtures of cross-linking ligands, such as antibodies, promote receptor internalization and degradation (30–32). To study the role of avidity in the behavior of phosphatidylserine-targeting PDCs, we also generated tetravalent Syt1-Fc-Syt1 that contains four Syt1 C2A domains (Fig. 2A). The tetravalent protein was purified as a homodimer (Fig. 2A; Supplementary Fig. S3A). As designed, Syt1-Fc-Syt1 has higher affinity/avidity for phosphatidylserine than Fc-Syt1 in lipid-binding assays (Fig. 2B) and the same lipid selectivity as its bivalent parent, Fc-Syt1 (Fig. 2C). Consistent with the binding data shown in Fig. 2B, the tetravalent Syt1-Fc-Syt1 bound to 2H11 cells at significantly higher levels than bivalent Fc-Syt1 protein (Fig. 2D).


We next compared the internalization of Fc-Syt1 and Syt1-Fc-Syt1 using Alexa 647–labeled proteins. 2H11 cells were incubated with labeled Fc-Syt1 and Syt1-Fc-Syt1 on ice at different concentrations to achieve similar surface binding followed by incubation at 37°C to allow internalization for different times. Surface-bound proteins were stripped off with EDTA (due to the Ca<sup>2+</sup>-dependent of binding), and internalized levels (resistant to stripping) were determined by flow cytometry. These studies indicated that although both proteins efficiently accumulated within cells, the tetravalent Syt1-Fc-Syt1 was internalized more rapidly compared with bivalent Fc-Syt1 (Fig. 2E). Fluorescent microscopy was also used to study the subcellular trafficking behavior of the Syt1-Fc fusion proteins. In both 2H11 and MDA-MB-231 cells, Fc-Syt1 and Syt1-Fc-Syt1 were internalized and delivered into LAMP-1 positive lysosomes (Fig. 2F and G; Supplementary Fig. S3B).

### Calcium sensing and endosomal release of phosphatidylserine-specific PDCs

The lysosomal trafficking and internalization behavior of Fc-Syt1 and Syt1-Fc-Syt1 indicated that they could be effective as

delivery vehicles for conjugated drugs. We therefore conjugated maleimidocaproyl-val-cit-PAB-MMAE (33) to hinge cysteines (Fig. 3A, left). For use as a negative control, MMAE was conjugated to a hen egg lysozyme-specific human IgG1 (21) in which the


light/heavy chain-interacting cysteine residues were mutated to serine to result in four possible conjugation sites. SDS-PAGE analyses indicated that the conjugations had gone to completion and resulted in drug-to-antibody ratios (DAR) of four (Fig. 3A,



**Figure 2.**

Cell binding and internalization of phosphatidylserine-specific Fc fusions containing Syt1 C2A. **A**, Schematic representation of bivalent and tetraivalent phosphatidylserine-specific Fc fusions with filled circles representing the Syt1 C2A domain (left). Right, reducing SDS-PAGE analyses of the Syt1-Fc fusions, with molecular weights (MW) shown in kDa on the right. **B**, Fc fusions (25 nmol/L) were incubated with nitrocellulose membranes coated with the indicated amounts of phosphatidylserine (PS). Bound proteins were detected with goat anti-human IgG (H + L) antibody conjugated with HRP. **C**, Lipid-binding profiles of Syt1-Fc fusions using lipid-coated nitrocellulose membranes as shown in Fig. 1B. Bound proteins were detected using goat anti-human IgG (H + L) antibody conjugated with HRP. **D**, 2H11 cells were treated with 50 nmol/L docetaxel for 72 hours, or treated with vehicle control (DMSO) and incubated with 50 nmol/L control IgG or phosphatidylserine-specific Fc fusions. Bound Fc fusion was detected using Alexa 488-labeled anti-human IgG (H + L), followed by flow cytometry analyses (MFI, mean fluorescence intensity). **E**, 2H11 cells were incubated with Alexa 647-labeled phosphatidylserine-specific Fc fusions on ice at optimized concentrations (220 nmol/L for Fc-Syt1 and 40 nmol/L for Syt1-Fc-Syt1) to achieve similar levels of surface binding. Cells were then incubated at 37°C for the indicated time periods. Surface-bound Fc fusions were stripped using 5 mmol/L EDTA and internalized proteins quantitated by flow cytometry analyses. **F** and **G**, 2H11 (F) or MDA-MB-231 (G) cells were incubated with 50 nmol/L control IgG or phosphatidylserine-specific Fc fusions at 37°C for 4 hours. Cells were fixed, stained with Cy3/Alexa 555-labeled anti-human IgG (H + L), and LAMP-1-specific antibody followed by Alexa 488-labeled secondary antibody for detecting LAMP-1. Fluorescence images were acquired and Cy3/Alexa 555, Alexa 488, and DAPI are pseudocolored red, green, and blue, respectively, in the overlays. Scale bar, 10 μm (F) and 5 μm (G). **D** and **E**, Statistically significant differences were analyzed using two-way ANOVA followed by Tukey *post hoc* test (\*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ ; \*\*\*\*,  $P < 0.0001$ ). Error bars in **D** and **E** represent SEM.

Li et al.

**Figure 3.**

Phosphatidylserine (PS)-specific PDCs bind to phosphatidylserine in a  $\text{Ca}^{2+}$ -dependent way and dissociate from phosphatidylserine in early endosomes prior to lysosomal delivery and disruption of microtubule networks in target cells. **A**, Schematic representation of phosphatidylserine-specific PDCs (left). The hinge cysteines were reduced and conjugated to four molecules of MMAE (small filled circles). Right, reducing SDS-PAGE analyses of the unconjugated or conjugated phosphatidylserine-specific Fc fusions and control IgG, with molecular weights (MW) shown in kDa on the left. **B**, phosphatidylserine-specific PDCs or MMAE-conjugated control IgG were incubated with phosphatidylserine-coated beads in the presence of the indicated  $\text{Ca}^{2+}$  concentrations (left) or pH levels (right). Bead-associated proteins were analyzed using immunoblotting and detection with goat anti-human IgG (H + L) conjugated with HRP. **C**, MDA-MB-231 cells were incubated with 100 nmol/L phosphatidylserine-specific PDCs or MMAE-conjugated control IgG for 30 minutes. Cells were fixed, stained with Alexa 555-labeled anti-human IgG (H + L), and EEA1-specific antibody followed by Alexa 488-labeled secondary antibody for detecting EEA1. Fluorescence images were acquired and Alexa 555, Alexa 488, and DAPI are pseudocolored red, green, and blue, respectively, in the overlays. The endosomes in boxed regions (labeled a and b) are cropped and expanded. Fluorescence intensities along the dotted lines in the overlays for these endosomes are shown in the fluorescence intensity plots. Scale bar, 5  $\mu\text{m}$  (left; whole cell images) and 1  $\mu\text{m}$  (right; cropped endosomes). **D**, MDA-MB-231 cells were incubated with 50 nmol/L MMAE-conjugated control IgG or phosphatidylserine-specific Fc fusions at 37°C for 4 hours. Cells were fixed, stained with Alexa 555-labeled anti-human IgG (H + L) and LAMP-1-specific antibody followed by Alexa 488-labeled secondary antibody for detecting LAMP-1. Fluorescence images were acquired and Alexa 555, Alexa 488, and DAPI are pseudocolored red, green, and blue, respectively, in the overlays. Scale bar, 10  $\mu\text{m}$ . **E** and **F**, 2H11 (**E**) and MDA-MB-231 (**F**) cells were treated with 100 or 50 nmol/L phosphatidylserine-specific PDCs or MMAE-conjugated control IgG for 10 or 20 hours, respectively. Cells were then fixed, stained with tubulin-specific antibody followed by Alexa 555-labeled secondary antibody, and imaged. Alexa 555 and DAPI are pseudocolored red and blue, respectively, in the overlays. Scale bar, 15  $\mu\text{m}$  (**E**) and 10  $\mu\text{m}$  (**F**).

right). The DARs of four for both MMAE-conjugated Fc-Syt1 and Syt1-Fc-Syt1 were confirmed by MALDI-TOF mass spectrometry. As complete conjugation disrupted the two disulfide bonds in the hinge region, apparent molecular weights of 43.6 kDa for Fc-Syt1\_MMAE and 58.6 kDa for Syt1-Fc-Syt1\_MMAE, respectively, were observed in these analyses (Supplementary Fig. S4A and S4B, top). This contrasted with the unconjugated or partially conjugated protein that retained two or one disulfide bonds, respectively, resulting in an apparent molecular weight of approximately 82 kDa for Fc-Syt1 and 112 kDa for Syt1-Fc-Syt1 (Supplementary Fig. S4A and S4B, bottom). Importantly, the conjugation process did not lead to protein aggregation (Supplementary Fig. S4C).

The Syt1 C2A domain requires Ca<sup>2+</sup> for phosphatidylserine binding (23). The lower Ca<sup>2+</sup> concentration in early/sorting endosomes (~2 μmol/L) compared with the extracellular Ca<sup>2+</sup> levels (1–2 mmol/L) suggests that following internalization, phosphatidylserine-specific PDCs will dissociate from the limiting membrane of these endosomes (18, 19). This dissociation is expected to lead to improved lysosomal delivery. Both phosphatidylserine-specific PDCs containing Syt1 domains bound to phosphatidylserine-beads in buffer containing 2 mmol/L Ca<sup>2+</sup>, but no detectable interaction was observed when the Ca<sup>2+</sup> concentration was decreased to 2 μmol/L (Fig. 3B, left). In addition, because the pH within sorting (early) endosomes is acidic (pH 6.0–6.5; ref. 34), we analyzed the effect of pH on PDC:PS interactions. Both PDCs bound to phosphatidylserine at similar levels in the pH range 6.0 to 7.4 (Fig. 3B, right). Consistent with the *in vitro* binding analyses demonstrating Ca<sup>2+</sup>-dependent binding, fluorescence microscopy analyses showed the presence of the phosphatidylserine-specific PDCs in the lumen, rather than limiting membrane of sorting endosomes in MDA-MB-231 cells following 15- and 30-minute incubation with PDC (Fig. 3C; Supplementary Fig. S5A). The relatively large size (~1.5 μm diameter) of the sorting endosomes in MDA-MB-231 cells, which was also observed with untreated cells (Supplementary Fig. S5B), enabled PDC dissociation from the limiting endosomal membrane to be quantitated (Fig. 3C, right). Fc-Syt1\_MMAE and Syt1-Fc-Syt1\_MMAE could be detected in lysosomes within 30 minutes, and the level of Fc fusion increased following 4 hours of incubation (Fig. 3D; Supplementary Fig. S5C). In addition, both PDCs disrupted the microtubular network in 2H11 and MDA-MB-231 cells (Fig. 3E and F).

#### Inhibition of growth and survival of phosphatidylserine-positive cells by phosphatidylserine-specific PDCs

We examined the effects of phosphatidylserine-specific PDCs on the growth of multiple cell lines, including tumor endothelium (2H11), ER-positive breast cancer (T-47D), HER2-positive breast cancer (SK-BR-3), triple-negative breast cancer (MDA-MB-231), androgen-sensitive prostate cancer (LNCaP), and castration-resistant prostate cancer (22Rv1). Staining of the cells with fluorescently labeled Annexin V showed that all of these cell lines were phosphatidylserine positive (Supplementary Fig. S6A). Incubation with phosphatidylserine-specific PDCs effectively inhibited the growth and survival of the cells in a dose-dependent manner (Fig. 4; Table 2). Although the tetravalent Syt1-Fc-Syt1\_MMAE was more potent in inhibiting T-47D cell growth than the bivalent Fc-Syt1\_MMAE, the two PDCs exhibited similar effects on the other cell lines. Consistent with the growth-inhibitory effects, the internalization of the two PDCs at concentrations close to the corresponding IC<sub>50</sub>s were similar for all cell lines,

except that Syt1-Fc-Syt1\_MMAE was internalized at higher levels than Fc-Syt1\_MMAE for T-47D cells (Supplementary Fig. S6B). Relatively high concentrations of the control IgG (hen egg lysozyme-specific human IgG1) conjugated with MMAE resulted in inhibition of cell growth, possibly due to nonspecific fluid phase uptake of the drug, whereas unconjugated phosphatidylserine-targeting proteins showed no effect on cell growth (Supplementary Fig. S6C). Thus, our data indicate that Syt1-based PDCs are potent inhibitors of tumor endothelial and cancer cell growth *in vitro*.

#### Inhibition of tumor growth by phosphatidylserine-specific PDCs in mouse xenograft models


We next investigated the therapeutic effects of the phosphatidylserine-specific PDCs against tumor xenografts in SCID mice. Prior to therapy, pharmacokinetic studies of the phosphatidylserine-specific PDCs demonstrated that tetravalent Syt1-Fc-Syt1\_MMAE had a shorter half-life than bivalent Fc-Syt1\_MMAE (Fig. 5A and B), possibly due to increased target-mediated uptake. Because of the relatively short half-lives, we delivered the PDCs (1 nmol/mouse) into tumor-bearing mice twice per week. In female SCID mice bearing orthotopic MDA-MB-231 breast tumors that had been pretreated with docetaxel, bivalent Fc-Syt1\_MMAE potentially blocked breast tumor growth (Fig. 5C). Tetravalent Syt1-Fc-Syt1\_MMAE also inhibited tumor growth, but was less effective than bivalent Fc-Syt1\_MMAE. Treatment with unconjugated Fc-Syt1 and Syt1-Fc-Syt1 had no effect on tumor growth (Fig. 5C).

In addition to efficacy in the breast tumor model, bivalent Fc-Syt1\_MMAE completely blocked tumor growth in male SCID mice bearing prostate cancer LNCaP xenografts that had been pretreated with docetaxel. Consistent with the *in vitro* data (Supplementary Fig. S6C), the unconjugated protein had no effect (Fig. 5D). However, tetravalent Syt1-Fc-Syt1\_MMAE did not significantly inhibit tumor growth. The most likely explanation for this difference is the shorter *in vivo* persistence of the tetravalent PDC. Taken together with the breast tumor data, these results indicate that the valency of the phosphatidylserine-specific PDCs has a profound influence on their therapeutic outcomes. Importantly, we did not observe weakness or loss of body weight in any of the treatment groups (Fig. 5E; Supplementary Fig. S7), indicating that phosphatidylserine-specific PDCs are well tolerated *in vivo*.

To further validate that Fc-Syt1\_MMAE binds phosphatidylserine-positive cells in tumor tissue following docetaxel treatment, we performed IHC following delivery of this PDC into tumor-bearing mice. We found that Fc-Syt1\_MMAE was localized to CD31-positive blood vessels (Fig. 5F), tumor cells, and tumor-infiltrating F4/80-positive macrophages (Fig. 5G), which can expose phosphatidylserine (5). The data indicate that cancer cells not only expose phosphatidylserine *in vitro* (Supplementary Fig. S6A; refs. 3, 35), but retain this loss of phosphatidylserine asymmetry *in vivo*.

#### Inhibition of tumor growth by Fc-Syt1\_MMAE is dependent on phosphatidylserine binding

To exclude the possibility that the drug accumulated in the tumor through nonspecific mechanisms, such as the enhanced permeability and retention effect (36), and to show that the *in vivo* efficacy was dependent on phosphatidylserine binding, we generated a mutated variant of the synaptotagmin 1 C2A domain with decreased affinity for phosphatidylserine. The C2A domain



**Figure 4.**

Phosphatidylserine-specific PDCs inhibit cancer cell growth and survival *in vitro*. Phosphatidylserine-positive cancer endothelial cells (2H11), breast cancer cells (T-47D, SK-BR-3 and MDA-MB-231), and prostate cancer cells (LNCaP and 22Rv1) were treated with antibody/protein-drug conjugates at the indicated concentrations. Cell viability following 72 hours (2H11), 96 hours (SK-BR-3, MDA-MB-231 and 22Rv1), or 120 hours (T-47D) is shown. Representative data from two or three independent experiments for each cell line are presented.

of synaptotagmin 1 interacts with phosphatidylserine through three  $\text{Ca}^{2+}$  ions chelated by five aspartic acids (D) in domain loops I and III (23). To ablate  $\text{Ca}^{2+}$  binding, we mutated all five aspartic acid residues to asparagines (N) to produce Fc-Syt1(DN). Fc-Syt1(DN) was purified as a homodimer and interacted with phosphatidylserine at essentially background levels in the protein–lipid overlay assay (Fig. 6A and B). Moreover, Fc-Syt1(DN) had significantly reduced binding to phosphatidylserine-positive cells compared with its wild-type counterpart (Fig. 6C). This mutated protein and as a control, human IgG1-derived Fc, were conjugated to MMAE and analyzed using mass spectrometry. These analyses indicated conjugation ratios of four drugs per protein molecule (Supplementary Fig. S8A and S8B). In addition, size exclusion analyses demonstrated that the resulting MMAE-conjugated proteins were not aggregated (Supplementary Fig. S8C). Mice bearing orthotopic MDA-MB-231 tumors were pre-treated with docetaxel prior to treatment with PDCs (1 nmol/mouse) or vehicle control for 4 weeks, until mice in the control (PBS) group were euthanized due to their large tumor sizes. Treatment of MDA-MB-231 tumors with Fc-Syt1\_MMAE led to potent growth inhibition (Fig. 6D). More importantly, tumor growth remained inhibited after the treatment with Fc-Syt1\_MMAE was stopped at 4 weeks. No tumors could be isolated from 3 of 6 mice at the end of the experiment (Fig. 6E). Although the delivery of Fc\_MMAE or Fc-Syt1(DN)\_MMAE slowed tumor growth initially, rapid proliferation was observed following the

end of treatment (Fig. 6D). Collectively, the data indicate that phosphatidylserine-binding is essential for the activity of Fc-Syt1\_MMAE.


## Discussion

ADCs represent a rapidly expanding class of therapeutics to target cancer. For the majority of ADC or PDC platforms, therapeutic efficacy is dependent on internalization and trafficking of the cytotoxic drug to lysosomes (11, 12). The current study describes a novel approach for the generation of a PDC to its target, namely exposed phosphatidylserine on tumor cells and tumor vascular endothelium. First, we have identified a phosphatidylserine-targeting protein that is efficiently internalized into cells that expose phosphatidylserine. Second, we have exploited the  $\text{Ca}^{2+}$  dependence of the phosphatidylserine–Syt1 interaction to enable endosomal dissociation, and consequent lysosomal delivery, of the Syt1-based PDC. Third, we have tuned the binding strength of the phosphatidylserine-targeting agent to optimize pharmacokinetic behavior and delivery to the tumor site. We demonstrate that the resulting PDC, Fc-Syt1\_MMAE, has potent antitumor effects in mouse models of breast and prostate cancer.


Among the three phosphatidylserine-targeting agents developed here, both Fc-Anx1 and Fc-PKC $\alpha$  exhibited higher levels of binding to target cells than Fc-Syt1. However, of these three agents, Fc-Syt1 exhibited the highest tumor accumulation in mice. The data indicate that high-affinity interactions with phosphatidylserine negatively impact the *in vivo* behavior with respect to tumor localization. There are two possible reasons for this behavior: First, phosphatidylserine exposure has also been identified in nonapoptotic normal cells, such as differentiating monocytes and

**Table 2.** IC<sub>50</sub> values for phosphatidylserine-specific PDCs

IC <sub>50</sub> (nmol/L)	2H11	T-47D	SK-BR-3	MDA-MB-231	LNCaP	22Rv1
Fc-Syt1_MMAE	7.0	14.1	1.2	9.1	7.6	6.0
Syt1-Fc-Syt1_MMAE	5.7	6.9	1.8	6.1	7.3	7.9

**Figure 5.**

Bivalent Fc-Syt1\_MMAE inhibits tumor growth *in vivo* and targets multiple phosphatidylserine-positive cells in the tumor tissue. **A**, Pharmacokinetic analyses of PS-PDCs in BALB/c SCID mice ( $n = 5$  mice/group). Whole body and blood radioactivities were measured at the indicated time points. **B**, AUCs in **A** were quantitated and statistically significant differences analyzed using unpaired Student *t* test (\*\*\*\*,  $P < 0.0001$ ). **C–E**, BALB/c SCID mice were implanted with MDA-MB-231 (**C** and **E**) or LNCaP (**D**) tumors. Mice ( $n = 5–6$  mice/group) were treated (day 33–60 in **C**, **E** and day 27–59 in **D**) with either unconjugated or MMAE-conjugated Fc fusions at a dose of 1 nmol/mouse (4.1 mg/kg for Fc-Syt1 or Fc-Syt1\_MMAE, 5.6 mg/kg for Syt1-Fc-Syt1 or Syt1-Fc-Syt1\_MMAE) twice per week. PBS was delivered as vehicle control. Tumor volumes (**C** and **D**) and body weights (**E**) were measured twice per week. Statistically significant differences (Fc-Syt1\_MMAE versus PBS in **C**; Fc-Syt1\_MMAE versus PBS or Syt1-Fc-Syt1\_MMAE in **D**) at treatment endpoints were analyzed using one-way ANOVA followed by Bonferroni *post hoc* test (\*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ ; \*\*\*\*,  $P < 0.0001$ ). Error bars in panels **A–E** represent SEM. **F** and **G**, BALB/c SCID mice bearing MDA-MB-231 tumors were treated (intraperitoneally) with 5 mg/kg docetaxel 72 and 48 hours before delivery of 1 nmol Fc-Syt1\_MMAE. PBS was delivered as vehicle control. Mice were perfused either 1 hour (**F**) or 24 hours (**G**) postinjection of Fc-Syt1\_MMAE. Tumors were dissected out, and tissue sections were fixed and stained with Alexa 555-labeled anti-human IgG (H + L), mouse CD31, F4/80, or human Ki-67-specific antibodies followed by Alexa 488-labeled secondary antibodies for detection of CD31/Ki67 and Alexa 647-labeled secondary antibody for detection of F4/80. Confocal images were acquired and Alexa 555, Alexa 488, Alexa 647, and DAPI are pseudocolored red, green, white, and blue, respectively, in the overlays. Scale bar, 50 μm (**F**) and 20 μm (**G**).

**Figure 6.**

Therapeutic effects of Fc-Syt1\_MMAE are dependent on phosphatidylserine binding. **A**, Reducing SDS-PAGE analyses of the unconjugated or MMAE-conjugated phosphatidylserine-specific Fc fusions and control IgG1 Fc, with molecular weights (MW) shown in kDa on the left. **B**, Lipid-coated nitrocellulose membranes were incubated with 2  $\mu$ g/mL Fc-Syt1 or Fc-Syt1(DN), and bound proteins detected using goat anti-human IgG antibody conjugated with HRP. **C**, 2H11 cells were treated with 50 nmol/L docetaxel for 72 hours, or treated with vehicle control (DMSO), and incubated with 5  $\mu$ g/mL control Fc, Fc-Syt1, or Fc-Syt1(DN). Bound Fc or Fc fusion was detected using Alexa 488-labeled anti-human IgG (H + L), followed by flow cytometry analyses. Statistically significant differences were analyzed using two-way ANOVA followed by Tukey *post hoc* test (\*\*\*,  $P < 0.001$ ; \*\*\*\*,  $P < 0.0001$ ). **D**, Female BALB/c SCID mice ( $n = 6$  mice/group) bearing MDA-MB-231 tumors were treated with the indicated agents at a dose of 1 nmol/mouse (4.1 mg/kg for Fc-Syt1\_MMAE or Fc-Syt1(DN)\_MMAE, 2.6 mg/kg for Fc\_MMAE) twice per week for 4 weeks (day 28–56), and tumor sizes were measured for a further 2.5 weeks (day 56–74). Statistically significant differences between Fc-Syt1\_MMAE and Fc-Syt1(DN)\_MMAE treatment groups at the treatment endpoint were analyzed using one-way ANOVA followed by Bonferroni *post hoc* test (\*\*\*,  $P < 0.001$ ). **E**, Tumors in each group shown in **D** were isolated and photographed. Scale bar, 1 cm. Error bars in **C** and **D** indicate SEM.

a subpopulation of T cells (16, 17). Although the SCID mice used in the current study lack T cells, monocytes may therefore act as a sink for target-mediated clearance outside the tumor tissue. This sink effect is expected to be greater for higher affinity phosphatidylserine-specific agents, resulting in shorter *in vivo* persistence. Consistent with this, pharmacokinetic studies demonstrate that Fc-Syt1 has a significantly longer half-life compared with Fc-AnxA1 and Fc-PKC $\alpha$ . Furthermore, bivalent Fc-Syt1\_MMAE also displayed a longer half-life and better *in vivo* efficacy than higher avidity, tetravalent Syt1-Fc-Syt1\_MMAE. Second, besides the influence on *in vivo* persistence, higher affinity HER2-specific antibodies have been shown to penetrate tumors less efficiently than their lower affinity counterparts [the "binding barrier theory" (37–39)]. Similar pathways may operate during phosphatidylserine targeting.

Interestingly, our study showed Fc-Syt1\_MMAE could potentially inhibit tumor growth in mice, whereas unconjugated Fc-Syt1 had no effect. This contrasts with bavituximab, which has been reported to be therapeutically effective as an unconjugated antibody in preclinical models when combined with radiation, chemotherapy, or other procedures that increase phosphatidylserine exposure in tumor tissues (6–10). One possible reason for this

difference is that we delivered the chemotherapeutic drug, docetaxel, as two doses at the beginning of the treatment to elevate tumor phosphatidylserine exposure, whereas in the bavituximab study, mice were treated in combination with antibody and docetaxel throughout therapy. The importance of increasing phosphatidylserine exposure using either chemotherapy or radiation for tumor localization of phosphatidylserine-targeting agents was indicated by a recent study in which two antibodies with  $\beta$ 2GP1-independent specificity for phosphatidylserine, phosphatidic acid, and cardiolipin failed to accumulate in tumor-bearing mice that were not treated with docetaxel or other "phosphatidylserine-inducing" agents (40).

The induction of endosomal dissociation of antibodies from their target antigen, by exploiting the lower pH or  $\text{Ca}^{2+}$  concentration within early/sorting endosomes relative to the extracellular environment, has been described by us and others as a strategy to clear antigens such as inflammatory cytokines or their soluble, signaling competent receptors (41–43). This release of target is followed by recycling and exocytosis of the antibody through binding to the Fc receptor, FcRn, which is expressed in most normal parenchymal and hematopoietic cells (20, 44, 45). To date, this approach has been implemented by using antibody

engineering to generate antibodies that bind with pH-dependence or Ca<sup>2+</sup>-dependence (41–43). Here, we use naturally occurring proteins that have intrinsic Ca<sup>2+</sup>-dependence for binding to target, thereby avoiding the need for engineering. Recent data including our own indicate that, in general, tumor cells express very low or undetectable levels of the recycling receptor FcRn (46, 47). This loss of expression is critical for efficacy of Ca<sup>2+</sup>-switched ADCs/PDCs, as it avoids the possibility of FcRn-mediated salvage of the drug conjugate away from lysosomal delivery following endosomal dissociation from target.

To summarize, we have developed a novel PDC that dissociates from target in early endosomes through Ca<sup>2+</sup> switching, enabling lysosomal delivery of the drug regardless of ligand (phosphatidylserine) trafficking. Our study also reveals a complex relationship between *in vivo* efficacy of the phosphatidylserine-targeting agent and *in vitro* binding properties, with the lowest affinity PDC, Fc-Syt1\_MMAE, showing superior pharmacokinetic properties and potent antitumor effects. Given that the subcellular trafficking behavior of many tumor surface markers following endosomal delivery remains poorly characterized, and most solid tumor antigens are not exclusively tumor specific, our study indicates that approaches involving Ca<sup>2+</sup> switching and affinity/avidity tuning may have value for the design of effective ADC or PDC platforms.

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Authors' Contributions

Conception and design: R. Li, L. Li, R.P. Mason, R.J. Ober, E.S. Ward

Development of methodology: R. Li, S. Chiguru, L. Li, R.P. Mason, E.S. Ward  
Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): R. Li, S. Chiguru, L. Li, S.C. Devanaboyina, H. Tian, A. Schroit, R.P. Mason, E.S. Ward

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): R. Li, S. Chiguru, D. Kim, R. Velmurugan, D. Kim, S.C. Devanaboyina, A. Schroit, R.P. Mason, R.J. Ober, E.S. Ward

Writing, review, and/or revision of the manuscript: R. Li, A. Schroit, R.P. Mason, R.J. Ober, E.S. Ward

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): R. Li, S. Chiguru, E.S. Ward

Study supervision: A. Schroit, R.J. Ober, E.S. Ward

## Acknowledgments

This study was supported in part by grants from the Cancer Prevention and Research Institute of Texas (CPRIT; RP 110441, awarded to E.S. Ward, R.J. Ober, A. Schroit, and R. Mason) and the NIH (Simmons Comprehensive Cancer Center support grant; 5P30CA142543, awarded to M. Cobb). The IVIS Spectrum was purchased with support (1S10RR024757, awarded to R.P. Mason) from the NIH.

This study is dedicated to the memory of our colleague, Philip Thorpe. We thank Alex Winters and Jeni Gerberich for assistance with the whole-body imaging experiments, Jacob Newman for assistance with the construction of expression plasmids for the phosphatidylserine-specific Fc fusions, and Drs. Dilip Challa and Priyanka Khare for providing recombinant mouse FcRn.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received January 30, 2017; revised May 4, 2017; accepted August 29, 2017; published OnlineFirst September 22, 2017.

## References

- Connor J, Bucana C, Fidler IJ, Schroit AJ. Differentiation-dependent expression of phosphatidylserine in mammalian plasma membranes: quantitative assessment of outer-leaflet lipid by prothrombinase complex formation. *Proc Natl Acad Sci U S A* 1989;86:3184–8.
- Ran S, Downes A, Thorpe PE. Increased exposure of anionic phospholipids on the surface of tumor blood vessels. *Cancer Res* 2002;62:6132–40.
- Riedl S, Rinner B, Asslaber M, Schaidt H, Walzer S, Novak A, et al. In search of a novel target - phosphatidylserine exposed by non-apoptotic tumor cells and metastases of malignancies with poor treatment efficacy. *Biochim Biophys Acta* 2011;1808:2638–45.
- Leventis PA, Grinstein S. The distribution and function of phosphatidylserine in cellular membranes. *Annu Rev Biophys* 2010;39:407–27.
- DeRose P, Thorpe PE, Gerber DE. Development of bavituximab, a vascular targeting agent with immune-modulating properties, for lung cancer treatment. *Immunotherapy* 2011;3:933–44.
- He J, Yin Y, Luster TA, Watkins L, Thorpe PE. Antiphosphatidylserine antibody combined with irradiation damages tumor blood vessels and induces tumor immunity in a rat model of glioblastoma. *Clin Cancer Res* 2009;15:6871–80.
- Yin Y, Huang X, Lynn KD, Thorpe PE. Phosphatidylserine-targeting antibody induces M1 macrophage polarization and promotes myeloid-derived suppressor cell differentiation. *Cancer Immunol Res* 2013;1:256–68.
- Huang X, Bennett M, Thorpe PE. A monoclonal antibody that binds anionic phospholipids on tumor blood vessels enhances the antitumor effect of docetaxel on human breast tumors in mice. *Cancer Res* 2005;65:4408–16.
- Cheng X, Li L, Thorpe PE, Yopp AC, Brekken RA, Huang X. Antibody-mediated blockade of phosphatidylserine enhances the antitumor effect of sorafenib in hepatocellular carcinomas xenografts. *Ann Surg Oncol* 2016;23:583–591.
- Beck AW, Luster TA, Miller AF, Holloway SE, Conner CR, Barnett CC, et al. Combination of a monoclonal anti-phosphatidylserine antibody with gemcitabine strongly inhibits the growth and metastasis of orthotopic pancreatic tumors in mice. *Int J Cancer* 2006;118:2639–43.
- Sievers EL, Senter PD. Antibody-drug conjugates in cancer therapy. *Annu Rev Med* 2013;64:15–29.
- Goldmacher VS, Kovtun YV. Antibody-drug conjugates: using monoclonal antibodies for delivery of cytotoxic payloads to cancer cells. *Ther Deliv* 2011;2:397–416.
- Bakhtiar R. Antibody drug conjugates. *Biotechnol Lett* 2016;38:1655–64.
- Chari RV, Miller ML, Widdison WC. Antibody-drug conjugates: an emerging concept in cancer therapy. *Angew Chem Int Ed Engl* 2014;53:3796–827.
- Kenis H, Reutlingsperger C. Targeting phosphatidylserine in anti-cancer therapy. *Curr Pharm Des* 2009;15:2719–23.
- Elliott JI, Surprenant A, Marelli-Berg FM, Cooper JC, Cassady-Cain RL, Wooding C, et al. Membrane phosphatidylserine distribution as a non-apoptotic signalling mechanism in lymphocytes. *Nat Cell Biol* 2005;7:808–16.
- Callahan MK, Halleck MS, Krahling S, Henderson AJ, Williamson P, Schlegel RA. Phosphatidylserine expression and phagocytosis of apoptotic thymocytes during differentiation of monocytic cells. *J Leukoc Biol* 2003;74:846–56.
- Albrecht T, Zhao Y, Nguyen TH, Campbell RE, Johnson JD. Fluorescent biosensors illuminate calcium levels within defined beta-cell endosome subpopulations. *Cell Calcium* 2015;57:263–74.
- Gerasimenko JV, Tepikin AV, Petersen OH, Gerasimenko OV. Calcium uptake via endocytosis with rapid release from acidifying endosomes. *Curr Biol* 1998;8:1335–8.
- Montoyo HP, Vaccaro C, Hafner M, Ober RJ, Mueller W, Ward ES. Conditional deletion of the MHC class I-related receptor FcRn reveals the sites of IgG homeostasis in mice. *Proc Natl Acad Sci U S A* 2009;106:2788–93.

Li et al.

21. Foote J, Winter G. Antibody framework residues affecting the conformation of the hypervariable loops. *J Mol Biol* 1992;224:487–99.
22. Neuberger MS. Expression and regulation of immunoglobulin heavy chain gene transfected into lymphoid cells. *EMBO J* 1983;2:1373–8.
23. Striegel AR, Biela LM, Evans CS, Wang Z, Delehoy JB, Sutton RB, et al. Calcium binding by synaptotagmin's C2A domain is an essential element of the electrostatic switch that triggers synchronous synaptic transmission. *J Neurosci* 2012;32:1253–60.
24. Popov S, Hubbard JG, Kim J, Ober B, Ghetie V, Ward ES. The stoichiometry and affinity of the interaction of murine Fc fragments with the MHC class I-related receptor, FcRn. *Mol Immunol* 1996;33:521–30.
25. Ober RJ, Ward ES. Compensation for loss of ligand activity in surface plasmon resonance experiments. *Anal Biochem* 2002;306:228–36.
26. Zhou J, Mateos F, Ober RJ, Ward ES. Conferring the binding properties of the mouse MHC class I-related receptor, FcRn, onto the human ortholog by sequential rounds of site-directed mutagenesis. *J Mol Biol* 2005;345:1071–81.
27. Schutters K, Reutelingsperger C. Phosphatidylserine targeting for diagnosis and treatment of human diseases. *Apoptosis* 2010;15:1072–82.
28. Gerke V, Creutz CE, Moss SE. Annexins: linking Ca<sup>2+</sup> signalling to membrane dynamics. *Nat Rev Mol Cell Biol* 2005;6:449–61.
29. Walter-Yohrling J, Morgenbesser S, Rouleau C, Bagley R, Callahan M, Weber W, et al. Murine endothelial cell lines as models of tumor endothelial cells. *Clin Cancer Res* 2004;10:2179–89.
30. Friedman LM, Rinon A, Schechter B, Lyass L, Lavi S, Bacus SS, et al. Synergistic down-regulation of receptor tyrosine kinases by combinations of mAbs: implications for cancer immunotherapy. *Proc Natl Acad Sci U S A* 2005;102:1915–20.
31. Spangler JB, Neil JR, Abramovitch S, Yarden Y, White FM, Lauffenburger DA, et al. Combination antibody treatment down-regulates epidermal growth factor receptor by inhibiting endosomal recycling. *Proc Natl Acad Sci U S A* 2010;107:13252–7.
32. Kang JC, Poovassery JS, Bansal P, You S, Manjarres IM, Ober RJ, et al. Engineering multivalent antibodies to target heregulin-induced HER3 signaling in breast cancer cells. *MAbs* 2014;6:340–53.
33. Doronina SO, Toki BE, Torgov MY, Mendelsohn BA, Cervený CG, Chace DF, et al. Development of potent monoclonal antibody auristatin conjugates for cancer therapy. *Nat Biotechnol* 2003;21:778–84.
34. Casey JR, Grinstein S, Orlowski J. Sensors and regulators of intracellular pH. *Nat Rev Mol Cell Biol* 2010;11:50–61.
35. Vallabhapurapu SD, Blanco VM, Sulaiman MK, Vallabhapurapu SL, Chu Z, Franco RS, et al. Variation in human cancer cell external phosphatidylserine is regulated by flippase activity and intracellular calcium. *Oncotarget* 2015;6:34375–88.
36. Matsumura Y, Maeda H. A new concept for macromolecular therapeutics in cancer chemotherapy: mechanism of tumorotropic accumulation of proteins and the antitumor agent smancs. *Cancer Res* 1986;46:6387–92.
37. Rudnick SI, Lou J, Shaller CC, Tang Y, Klein-Szanto AJ, Weiner LM, et al. Influence of affinity and antigen internalization on the uptake and penetration of anti-HER2 antibodies in solid tumors. *Cancer Res* 2011;71:2250–9.
38. Thurber GM, Schmidt MM, Wittrup KD. Antibody tumor penetration: transport opposed by systemic and antigen-mediated clearance. *Adv Drug Deliv Rev* 2008;60:1421–34.
39. Adams GP, Schier R, McCall AM, Simmons HH, Horak EM, Alpaugh RK, et al. High affinity restricts the localization and tumor penetration of single-chain Fv antibody molecules. *Cancer Res* 2001;61:4750–5.
40. Bujak E, Pretto F, Neri D. Generation and tumor recognition properties of two human monoclonal antibodies specific to cell surface anionic phospholipids. *Invest New Drugs* 2015;33:791–800.
41. Igawa T, Ishii S, Tachibana T, Maeda A, Higuchi Y, Shimaoka S, et al. Antibody recycling by engineered pH-dependent antigen binding improves the duration of antigen neutralization. *Nat Biotechnol* 2010;28:1203–7.
42. Hironiwa N, Ishii S, Kadono S, Iwayanagi Y, Mimoto F, Habu K, et al. Calcium-dependent antigen binding as a novel modality for antibody recycling by endosomal antigen dissociation. *MAbs* 2016;8:65–73.
43. Devanaboyina SC, Lynch SM, Ober RJ, Ram S, Kim D, Puig-Canto A, et al. The effect of pH dependence of antibody-antigen interactions on subcellular trafficking dynamics. *MAbs* 2013;5:851–9.
44. Zhu X, Meng G, Dickinson BL, Li X, Mizoguchi E, Miao L, et al. MHC class I-related neonatal Fc receptor for IgG is functionally expressed in monocytes, intestinal macrophages, and dendritic cells. *J Immunol* 2001;166:3266–76.
45. Akilesh S, Christianson GJ, Roopenian DC, Shaw AS. Neonatal FcR expression in bone marrow-derived cells functions to protect serum IgG from catabolism. *J Immunol* 2007;179:4580–8.
46. Swiercz R, Chiguru S, Tahmasbi A, Ramezani SM, Hao G, Challa DK, et al. Use of Fc-engineered antibodies as clearing agents to increase contrast during PET. *J Nucl Med* 2014;55:1204–7.
47. Dalloneau E, Baroukh N, Mavridis K, Maillet A, Gueugnon F, Courty Y, et al. Downregulation of the neonatal Fc receptor expression in non-small cell lung cancer tissue is associated with a poor prognosis. *Oncotarget* 2016;7:54415–29.

# Molecular Cancer Therapeutics

## Targeting Phosphatidylserine with Calcium-Dependent Protein–Drug Conjugates for the Treatment of Cancer

Ran Li, Srinivas Chiguru, Li Li, et al.

*Mol Cancer Ther* 2018;17:169-182. Published OnlineFirst September 22, 2017.

**Updated version** Access the most recent version of this article at:  
doi:[10.1158/1535-7163.MCT-17-0092](https://doi.org/10.1158/1535-7163.MCT-17-0092)

**Supplementary Material** Access the most recent supplemental material at:  
<http://mct.aacrjournals.org/content/suppl/2017/09/22/1535-7163.MCT-17-0092.DC1>

**Cited articles** This article cites 47 articles, 16 of which you can access for free at:  
<http://mct.aacrjournals.org/content/17/1/169.full#ref-list-1>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://mct.aacrjournals.org/content/17/1/169>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.